



07/17/97

A/Sequence

PATENT

Docket No. P50533-03

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Martin Karl Russel Burnham, et al.

For: NOVEL XANTHINE PHOSPHORIBOSYL TRANSFERASE

Assistant Commissioner for Patents
Washington, D.C. 20231**COVER SHEET FOR FILING PATENT APPLICATION**

Enclosed herewith for filing in the United States Patent and Trademark Office are the Specification and Declaration and Power of Attorney for the patent application identified below:

1. The names of the inventors are: Martin Karl Russel Burnham, Michael Terrence Black, John Edward Hodgson, David Justin Charles Knowles, Michael Arthur Lonetto, Richard Oakley Nicholas, and Robert King Stodola.
2. The addresses of the inventors are:
Martin Karl Russel Burnham
2927 Tanglewood Lane
Norristown, Pennsylvania, USA 19403
(A Citizen of the United Kingdom)

Michael Terrence Black
502 Milhouse Way
Chester Springs, Pennsylvania, USA 19425
(A Citizen of the United Kingdom)

John Edward Hodgson
260 Lapp Road
Malvern, Pennsylvania, USA 19355
(A Citizen of the United Kingdom)

2. The addresses of the inventors (continued):
- David Justin Charles Knowles
Downsview House, 45 Cronks Hill Road
Redhill, Surrey, United Kingdom RH1 6LY
(A Citizen of the United Kingdom)
- Michael Arthur Lonetto
18 Victoria Circle
Collegeville, Pennsylvania, USA 19426
(A Citizen of the United States)
- Richard Oakly Nicholas
355 Carmen Drive
Collegeville, Pennsylvania, USA 19426
(A Citizen of the United Kingdom)
- Robert King Stodola
309 Haws Lane
Flourtown, Pennsylvania, USA 19031
(A Citizen of the United States)
3. The title of the invention is: Novel Xanthine Phosphoribosyl Transferase
4. The name, registration and telephone number of the attorney is:
- Attorney: Arthur E. Jackson
Registration No. 34,354
Telephone No. 609/520-3254
5. The Docket Number used to identify this application is: P50533-03
6. The correspondence address for this application is:
- Q. Todd Dickinson
Dechert, Price & Rhoads
4000 Bell Atlantic Tower, 1717 Arch Street
Philadelphia, PA 19103-2793


This invention was not made by an agency of the United States Government, or under contract with an agency of the United States Government.

This application claims benefit of United States application number 60/024,022, filed August 16, 1996.

There are 40 pages of specification, 0 sheets of drawings and 20 claims accompanying this cover sheet.

The filing fee for this patent application, is \$770.00. Please charge the \$770.00 patent application fee, and any deficiencies to this fee, to Deposit Account No. 04-0475. A duplicate copy of this cover sheet is enclosed for accounting purposes.

If the fee required for this application is different from the amount identified above, the proper amount should be charged to Deposit Account No. 04-0475.




Stephen T. Falk
Registration No. 36,795
Dechert, Price & Rhoads
4000 Bell Atlantic Tower
1717 Arch Street
Philadelphia, PA 19103-2793, USA
Phone: 215/994-2488

Date: July 17, 1997

CERTIFICATION UNDER 37 CFR 1.10

I hereby certify that this correspondence and the documents referred to as attached therein are being deposited with the United States Postal Service on **July 17, 1997**, in an envelope as "EXPRESS MAIL POST OFFICE TO ADDRESSEE" service under 37 CFR 1.10, Mailing Label Number **EI 522 227 725 US** addressed to: Assistant Commissioner for Patents, Washington, D.C. 20231.


Mary Katherine Morgan Merlino

INVENTORS:

MARTIN KARL RUSSEL BURNHAM
2927 Tanglewood Lane
Norristown, Pennsylvania, USA 19403
(A Citizen of the United Kingdom)

MICHAEL TERENCE BLACK
502 Milhouse Way
Chester Springs, Pennsylvania, USA 19425
(A Citizen of the United Kingdom)

JOHN EDWARD HODGSON
260 Lapp Road
Malvern, Pennsylvania, USA 19355
(A Citizen of the United Kingdom)

DAVID JUSTIN CHARLES KNOWLES
Downsview House, 45 Cronks Hill Road
Redhill, Surrey, United Kingdom RH1 6LY
(A Citizen of the United Kingdom)

MICHAEL ARTHUR LONETTO
18 Victoria Circle
Collegeville, Pennsylvania, USA 19426
(A Citizen of the United States)

RICHARD OAKLY NICHOLAS
355 Carmen Drive
Collegeville, Pennsylvania, USA 19426
(A Citizen of the United Kingdom)

ROBERT KING STODOLA
309 Haws Lane
Flourtown, Pennsylvania, USA 19031
(A Citizen of the United States)

TITLE:

**NOVEL XANTHINE PHOSPHORIBOSYL
TRANSFERASE**

NOVEL XANTHINE PHOSPHORIBOSYL TRANSFERASE

RELATED APPLICATIONS

5 This application claims benefit of United States application number 60/024,022, filed August 16, 1996.

FIELD OF THE INVENTION

10 This invention relates to newly identified polynucleotides and polypeptides, and their production and uses, as well as their variants, agonists and antagonists, and their uses. In particular, in these and in other regards, the invention relates to novel polynucleotides and polypeptides of the genes associated with purine salvage family, hereinafter referred to as "xanthine phosphoribosyl transferase".

15

BACKGROUND OF THE INVENTION

 The Streptococci make up a medically important genera of microbes known to cause several types of disease in humans, including, for example, otitis media, conjunctivitis, pneumonia, bacteremia, meningitis, sinusitis, pleural empyema and endocarditis, and most particularly meningitis, such as for example infection of cerebrospinal fluid. Since its isolation more than 100 years ago, *Streptococcus pneumoniae* has been one of the more intensively studied microbes. For example, much of our early understanding that DNA is, in fact, the genetic material was predicated on the work of Griffith and of Avery, Macleod and McCarty using this microbe. Despite the vast amount of research with *S. pneumoniae*, many questions concerning the virulence of this microbe remain. It is particularly preferred to employ Streptococcal genes and gene products as targets for the development of antibiotics.

 The frequency of *Streptococcus pneumoniae* infections has risen dramatically in the past 20 years. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to

isolate *Streptococcus pneumoniae* strains which are resistant to some or all of the standard antibiotics. This has created a demand for both new anti-microbial agents and diagnostic tests for this organism.

Xanthine phosphoribosyl transferase is an enzyme involved in purine salvage under
5 nitrogen limiting conditions such as might be prevalent during some stages of disease caused by pathogens containing this pathway (see Christiansen LC, Schou S, Nygaard P, Saxild HH J Bacteriol 179 (8): 2540-2550 (Apr 1997)) Xanthine metabolism in *Bacillus subtilis*: characterization of the xpt-pbuX operon and evidence for purine- and nitrogen-controlled expression of genes involved in xanthine salvage and catabolism.)

10 Clearly, there is a need for factors, such as the novel compounds of the invention, that have a present benefit of being useful to screen compounds for antibiotic activity. Such factors are also useful to determine their role in pathogenesis of infection, dysfunction and disease. There is also a need for identification and characterization of such factors and their antagonists and agonists which can play a role in preventing, ameliorating or correcting infections, dysfunctions
15 or diseases.

The polypeptides of the invention have amino acid sequence homology to a known the xanthine phosphoribosyl transferase of *B.subtilis* (XPT_BACSU) protein.

SUMMARY OF THE INVENTION

20 It is an object of the invention to provide polypeptides that have been identified as novel xanthine phosphoribosyl transferase polypeptides by homology between the amino acid sequence set out in Table 1 [SEQ ID NO: 2] and a known amino acid sequence or sequences of other proteins such as the xanthine phosphoribosyl transferase of *B.subtilis* (XPT_BACSU) protein.

25 It is a further object of the invention to provide polynucleotides that encode xanthine phosphoribosyl transferase polypeptides, particularly polynucleotides that encode the polypeptide herein designated xanthine phosphoribosyl transferase.

In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding xanthine phosphoribosyl transferase polypeptides comprising the sequence set
30 out in Table 1 [SEQ ID NO:1] which includes a full length gene, or a variant thereof.

In another particularly preferred embodiment of the invention there is a novel xanthine phosphoribosyl transferase protein from *Streptococcus pneumoniae* comprising the amino acid sequence of Table 1 [SEQ ID NO:2], or a variant thereof.

5 In accordance with another aspect of the invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the *Streptococcus pneumoniae* 0100993 strain contained in the deposited strain.

A further aspect of the invention there are provided isolated nucleic acid molecules encoding xanthine phosphoribosyl transferase, particularly *Streptococcus pneumoniae* xanthine phosphoribosyl transferase, including mRNAs, cDNAs, genomic DNAs. Further embodiments of
10 the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

In accordance with another aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization. Among the particularly preferred embodiments of the invention are naturally
15 occurring allelic variants of xanthine phosphoribosyl transferase and polypeptides encoded thereby.

Another aspect of the invention there are provided novel polypeptides of *Streptococcus pneumoniae* referred to herein as xanthine phosphoribosyl transferase as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and
20 compositions comprising the same.

Among the particularly preferred embodiments of the invention are variants of xanthine phosphoribosyl transferase polypeptide encoded by naturally occurring alleles of the xanthine phosphoribosyl transferase gene.

In a preferred embodiment of the invention there are provided methods for producing the
25 aforementioned xanthine phosphoribosyl transferase polypeptides.

In accordance with yet another aspect of the invention, there are provided inhibitors to such polypeptides, useful as antibacterial agents, including, for example, antibodies.

In accordance with certain preferred embodiments of the invention, there are provided products, compositions and methods for assessing xanthine phosphoribosyl transferase
30 expression, treating disease, for example, otitis media, conjunctivitis, pneumonia, bacteremia, meningitis, sinusitis, pleural empyema and endocarditis, and most particularly meningitis, such as

for example infection of cerebrospinal fluid, assaying genetic variation, and administering a xanthine phosphoribosyl transferase polypeptide or polynucleotide to an organism to raise an immunological response against a bacteria, especially a *Streptococcus pneumoniae* bacteria.

5 In accordance with certain preferred embodiments of this and other aspects of the invention there are provided polynucleotides that hybridize to xanthine phosphoribosyl transferase polynucleotide sequences, particularly under stringent conditions.

In certain preferred embodiments of the invention there are provided antibodies against xanthine phosphoribosyl transferase polypeptides.

10 In other embodiments of the invention there are provided methods for identifying compounds which bind to or otherwise interact with and inhibit or activate an activity of a polypeptide or polynucleotide of the invention comprising: contacting a polypeptide or polynucleotide of the invention with a compound to be screened under conditions to permit binding to or other interaction between the compound and the polypeptide or polynucleotide to assess the binding to or other interaction with the compound, such binding or interaction being
15 associated with a second component capable of providing a detectable signal in response to the binding or interaction of the polypeptide or polynucleotide with the compound; and determining whether the compound binds to or otherwise interacts with and activates or inhibits an activity of the polypeptide or polynucleotide by detecting the presence or absence of a signal generated from the binding or interaction of the compound with the polypeptide or polynucleotide.

20 In accordance with yet another aspect of the invention, there are provided xanthine phosphoribosyl transferase agonists and antagonists, preferably bacteriostatic or bacteriocidal agonists and antagonists.

In a further aspect of the invention there are provided compositions comprising a xanthine phosphoribosyl transferase polynucleotide or a xanthine phosphoribosyl transferase
25 polypeptide for administration to a cell or to a multicellular organism.

Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

30

GLOSSARY

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Host cell" is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

5 "Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods,
10 including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two
20 sequences include, but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Atschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990)). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990)). As an illustration, by a
25 polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having a
30 nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or

a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. Analogously, by a polypeptide having an amino acid sequence having at least, for example, 95% identity to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded regions, or a mixture of single- and double-stranded regions. In addition, "polynucleotide" as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different

molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with
5 backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term
10 "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including, for example, simple and complex cells. "Polynucleotide(s)" also embraces short polynucleotides often referred to as oligonucleotide(s).

"Polypeptide(s)" refers to any peptide or protein comprising two or more amino acids
15 joined to each other by peptide bonds or modified peptide bonds. "Polypeptide(s)" refers to both short chains, commonly referred to as peptides, oligopeptides and oligomers and to longer chains generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such as processing and other post-translational modifications, but also by chemical modification
20 techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. It will be appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Modifications can occur anywhere in a polypeptide,
25 including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Modifications include, for example, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation,
30 formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination,

methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, 5 *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993) and Wold, F., *Posttranslational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter et al., *Meth. Enzymol.* 182:626-646 (1990) and Rattan et al., *Protein Synthesis: Posttranslational* 10 *Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

"Variant(s)" as the term is used herein, is a polynucleotide or polypeptide that differs 15 from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in 20 the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any 25 combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques, by direct synthesis, and by other recombinant methods known to skilled artisans.

30

DESCRIPTION OF THE INVENTION

The invention relates to novel xanthine phosphoribosyl transferase polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of a novel xanthine phosphoribosyl transferase of *Streptococcus pneumoniae*, which is related by amino acid sequence homology to the xanthine phosphoribosyl transferase of *B.subtilis* (XPT_BACSU) polypeptide. The invention relates especially to xanthine phosphoribosyl transferase having the nucleotide and amino acid sequences set out in Table 1 [SEQ ID NO: 1] and Table 1 [SEQ ID NO: 2] respectively, and to the xanthine phosphoribosyl transferase nucleotide sequences of the DNA in the deposited strain and amino acid sequences encoded thereby.

TABLE 1
xanthine phosphoribosyl transferase Polynucleotide and Polypeptide Sequences

(A) Sequences from *Streptococcus pneumoniae* xanthine phosphoribosyl transferase polynucleotide sequence [SEQ ID NO:1].

5'-1 ATGAAATTAT TAGAAGAGCG CATCCTCAAG GATGGGCATA TCTTGGGTGA

51 TAACATCCTC AAGGTAGATT CCTTTTAAAC CCACCAAGTT GACTTTAGCT

101 TGATGCGAGA GATTGGTAAG GTTTTTGCGG AAAAATTTGC TGCTACTGGC

151 ATTACCAAGG TCGTAACCAT TGAAGCGTCG GGTATTGCCC CAGCCGTTTT

201 TACAGCTGAA GCCTTAAACG TTCCCATGAT TTTCGCCAAA AAAGCTAAGA

251 ACATCACCAT GAACGAAGGC ATCTTAACTG CTCAAGTCTA CTCCTTTACC

301 AAGCAGGTGA CCAGCACTGT TTCTATCGCT GGAAAATTCC TCTCACCAGA

351 GGACAAGGTT TTGATTATCG ACGATTTCTT TGCTAATGGC CAAGCTGCTA

401 AAGGCTTGAT TCAAATCATC GAACAGGCCG GTGCCACAGT CCAAGCTATC

451 GGTATCGTGA TTGAGAAATC CTTCCAAGAT GGTCGTGATT TGCTTGAAAA

5 501 AGCAGGCTAC CCTGTCCTAT CACTTGCTCG CTTGGATCGT TTTGAAAATG

551 GTCAGGTCGT ATTTAAGGAG GCAGATCTCT AA-3'

(B) xanthine phosphoribosyl transferase polypeptide sequence deduced from the
10 polynucleotide sequence in this table [SEQ ID NO:2].

NH₂-1 MKLLEERILK DGHILGDNIL KVDSFLTHQV DFSLMREIGK VFAEKFAATG

51 ITKVVITIEAS GIAPAVFTAE ALNVP MIFAK KAKNITMNEG ILTAQVYSFT

15 101 KQVTSTVSIA GKFLSPEDKV LIIDDFLANG QAAKGLIQII EQAGATVQAI

151 GIVIEKSFQD GRDLLEKAGY PVLSLARLDR FENGQVVFKE ADL*-COOH

(C) Polynucleotide sequence embodiments [SEQ ID NO:1].

20 X-(R₁)_n-1 ATGAAATTAT TAGAAGAGCG CATCCTCAAG GATGGGCATA TCTTGGGTGA

51 TAACATCCTC AAGGTAGATT CCTTTTAAAC CCACCAAGTT GACTTTAGCT

101 TGATGCGAGA GATTGGTAAG GTTTTTGCGG AAAAATTTGC TGCTACTGGC

25 151 ATTACCAAGG TCGTAACCAT TGAAGCGTCG GGTATTGCCC CAGCCGTTTT

201 TACAGCTGAA GCCTTAAACG TTCCCATGAT TTTCGCCAAA AAAGCTAAGA

30 251 ACATCACCAT GAACGAAGGC ATCTTAACTG CTCAAGTCTA CTCCTTTACC

301 AAGCAGGTGA CCAGCACTGT TTCTATCGCT GGAAAATTCC TCTCACCAGA

351 GGACAAGGTT TTGATTATCG ACGATTTCTT TGCTAATGGC CAAGCTGCTA

401 AAGGCTTGAT TCAAATCATC GAACAGGCCG GTGCCACAGT CCAAGCTATC

5 451 GGTATCGTGA TTGAGAAATC CTTCCAAGAT GGTCTGTGATT TGCTTGAAAA

501 AGCAGGCTAC CCTGTCCTAT CACTTGCTCG CTTGGATCGT TTTGAAAATG

551 GTCAGGTCGT ATTTAAGGAG GCAGATCTCT AA- (R₂)_n-Y

10 (D) Polypeptide sequence embodiments [SEQ ID NO:2].

X- (R₁)_n-1 MKLLEERILK DGHILGDNIL KVDSFLTHQV DFSLMREIGK VFAEKFAATG

51 ITKVVTIEAS GIAPAVFTAE ALNVP MIFAK KAKNITMNEG ILTAQVYSFT

15 101 KQVTSTVSIA GKFLSPEDKV LIIDDFLANG QAAKGLIQII EQAGATVQAI

151 GIVIEKSFQD GRDLLEKAGY PVL SLARLDR FENGQVVFKE ADL* - (R₂)_n-Y

20 (E) Sequences from Streptococcus pneumoniae xanthine phosphoribosyl transferase polynucleotide ORF sequence [SEQ ID NO:3].

5'-1 ATGAAATTAT TAGAAGAGCG CATCCTCAAG GATGGGCATA TCTTGGGTGA

51 TAACATCCTC AAGGTAGATT CCTTTTAAAC CCACCAAGTT GACTTTAGCT

25 101 TGATGCGAGA GATTGGTAAG GTTTTTCGGG AAAAATTGTC TGCTACTGGC

151 ATTACCAAGG TCGTAACCAT TGAAGCGTCG GGTATTGCCC CAGCCGTTTT

30 201 TACAGCTGAA GCCTTAAACG TTCCCATGAT TTTCGCCAAA AAAGCTAAGA

251 ACATCACCAT GAACGAAGGC ATCTTAACTG CTCAAGTCTA CTCCTTTACC

301 AAGCAGGTGA CCAGCACTGT TTCTATCGCT GGAAAATTCC TCTCACCAGA
 351 GGACAAGGTT TTGATTATCG ACGATTTCCT TGCTAATGGC CAAGCTGCTA
 5 401 AAGGCTTGAT TCAAATCATC GAACAGGCCG GTGCCACAGT CCAAGCTATC
 451 GGTATCGTGA TTGAGAAATC CTTCCAAGAT GGTCGTGATT TGCTTGAAAA
 501 AGCAGGCTAC CCTGTCCTAT CACTTGCTCG CTTGGATCGT TTTGAAAATG
 10 551 GTCAGGTCGT ATTTAAGGAG GCAGATCTC-3'

(F) xanthine phosphoribosyl transferase polypeptide sequence deduced from the
 polynucleotide ORF sequence in this table [SEQ ID NO:4].
 15 NH₂-MKLLEERILK DGHILGDNIL KVDSFLTHQV DFSLMREIGK VFAEKFAATG

51 ITKVVTIEAS GIAPAVFTAE ALNVP MIFAK KAKNITMNEG ILTAQVYSFT
 101 KQVTSTVSIA GKFLSPEDKV LIIDDFLANG QAAKGLIQII EQAGATVQAI
 20 151 GIVIEKSFQD GRDLLEKAGY PVLSLARLDR FENGQVVFKE ADL-COOH

Deposited materials

A deposit containing a *Streptococcus pneumoniae* 0100993 strain has been deposited
 25 with the National Collections of Industrial and Marine Bacteria Ltd. (herein "NCIMB"), 23 St.
 Machar Drive, Aberdeen AB2 1RY, Scotland on 11 April 1996 and assigned deposit number
 40794. The deposit was described as *Streptococcus peumnoiae* 0100993 on deposit. On 17 April
 1996 a *Streptococcus peumnoiae* 0100993 DNA library in *E. coli* was similarly deposited with the
 NCIMB and assigned deposit number 40800.. The *Streptococcus pneumoniae* strain deposit is
 30 referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

The deposited strain contains the full length xanthine phosphoribosyl transferase gene.
 The sequence of the polynucleotides contained in the deposited strain, as well as the amino acid

sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

The deposit of the deposited strain has been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent
5 Procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strain is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.

A license may be required to make, use or sell the deposited strain, and compounds
10 derived therefrom, and no such license is hereby granted.

Polypeptides

The polypeptides of the invention include the polypeptide of Table 1 [SEQ ID NO:2] (in particular the mature polypeptide) as well as polypeptides and fragments, particularly those which have the biological activity of xanthine phosphoribosyl transferase, and also those which have at
15 least 70% identity to a polypeptide of Table 1 [SEQ ID NOS:2 and 4] or the relevant portion, preferably at least 80% identity to a polypeptide of Table 1 [SEQ ID NOS:2 and 4], and more preferably at least 90% similarity (more preferably at least 90% identity) to a polypeptide of Table 1 [SEQ ID NOS:2 and 4] and still more preferably at least 95% similarity (still more preferably at least 95% identity) to a polypeptide of Table 1 [SEQ ID NOS:2 and 4] and also
20 include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

The invention also includes polypeptides of the formula set forth in Table 1 (D) [SEQ ID NO:2] wherein, at the amino terminus, X is hydrogen, and at the carboxyl terminus, Y is hydrogen or a metal, R₁ and R₂ is any amino acid residue, and n is an integer between 1 and
25 1000. Any stretch of amino acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

A fragment is a variant polypeptide having an amino acid sequence that entirely is the same as part but not all of the amino acid sequence of the aforementioned polypeptides. As with xanthine phosphoribosyl transferase polypeptides fragments may be "free-standing," or comprised
30 within a larger polypeptide of which they form a part or region, most preferably as a single continuous region, a single larger polypeptide.

Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of Table 1 [SEQ ID NOS:2 and 4], or of variants thereof, such as a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus. Degradation forms of the polypeptides of the invention in a host cell, particularly a *Streptococcus pneumoniae*, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

Also preferred are biologically active fragments which are those fragments that mediate activities of xanthine phosphoribosyl transferase, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those fragments that are antigenic or immunogenic in an animal, especially in a human. Particularly preferred are fragments comprising receptors or domains of enzymes that confer a function essential for viability of *Streptococcus pneumoniae* or the ability to initiate, or maintain cause disease in an individual, particularly a human.

Variants that are fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention.

Polynucleotides

Another aspect of the invention relates to isolated polynucleotides, including the full length gene, that encode the xanthine phosphoribosyl transferase polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NOS:2 and 4] and polynucleotides closely related thereto and variants thereof.

Using the information provided herein, such as a polynucleotide sequence set out in Table 1 [SEQ ID NOS:1 and 3], a polynucleotide of the invention encoding xanthine phosphoribosyl transferase polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Streptococcus pneumoniae* 0100993 cells as starting material, followed by obtaining a full length

clone. For example, to obtain a polynucleotide sequence of the invention, such as a sequence given in Table 1 [SEQ ID NOS:1 and 3], typically a library of clones of chromosomal DNA of *Streptococcus pneumoniae* 0100993 in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence.

5 Clones carrying DNA identical to that of the probe can then be distinguished using stringent conditions. By sequencing the individual clones thus identified with sequencing primers designed from the original sequence it is then possible to extend the sequence in both directions to determine the full gene sequence. Conveniently, such sequencing is performed using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are

10 described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Illustrative of the invention, the polynucleotide set out in Table 1 [SEQ ID NO:1] was discovered in a DNA library derived from

15 *Streptococcus pneumoniae* 0100993.

The DNA sequence set out in Table 1 [SEQ ID NOS:1] contains an open reading frame encoding a protein having about the number of amino acid residues set forth in Table 1 [SEQ ID NO:2] with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known in the art. The polynucleotide of SEQ ID NO: 1, between

20 nucleotide number 1 through number 579 encodes the polypeptide of SEQ ID NO:2. The stop codon begins at nucleotide number 582 of SEQ ID NO:1.1579

Xanthine phosphoribosyl transferase of the invention is structurally related to other proteins of the genes associated with purine salvage family, as shown by the results of sequencing the DNA encoding xanthine phosphoribosyl transferase of the deposited strain. The

25 protein exhibits greatest homology to the xanthine phosphoribosyl transferase of *B.subtilis* (XPT_BACSU) protein among known proteins. Xanthine phosphoribosyl transferase of Table 1 [SEQ ID NO:2] has about 58%% identity over its entire length and about 75%% similarity over its entire length with the amino acid sequence of the xanthine phosphoribosyl transferase of *B.subtilis* (XPT_BACSU) polypeptide.

30 The invention provides a polynucleotide sequence identical over its entire length to the coding sequence in Table 1 [SEQ ID NO:1]. Also provided by the invention is the coding

sequence for the mature polypeptide or a fragment thereof, by itself as well as the coding sequence for the mature polypeptide or a fragment in reading frame with other coding sequence, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence. The polynucleotide may also contain non-coding sequences, including for example, but not
5 limited to non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals, and additional coding sequence which encode additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine
10 peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA tag (Wilson *et al.*, *Cell* 37: 767 (1984). Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

A preferred embodiment of the invention is a polynucleotide of comprising nucleotide 1
15 to 579 or 582 set forth in SEQ ID NO:1 of Table 1 which encode the xanthine phosphoribosyl transferase polypeptide.

The invention also includes polynucleotides of the formula set forth in Table 1 (C)[SEQ ID NO:1] wherein, at the 5' end of the molecule, X is hydrogen, and at the 3' end of the molecule, Y is hydrogen or a metal, R₁ and R₂ is any nucleic acid residue, and n is an integer between 1 and
20 1000. Any stretch of nucleic acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Streptococcus pneumoniae*
25 xanthine phosphoribosyl transferase having the amino acid sequence set out in Table 1 [SEQ ID NO:2]. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, interrupted by integrated phage or an insertion sequence or editing) together with additional regions, that also may contain coding and/or non-coding sequences.

30 The invention further relates to variants of the polynucleotides described herein that encode for variants of the polypeptide having the deduced amino acid sequence of Table 1 [SEQ

ID NO:2]. Variants that are fragments of the polynucleotides of the invention may be used to synthesize full-length polynucleotides of the invention.

Further particularly preferred embodiments are polynucleotides encoding xanthine phosphoribosyl transferase variants, that have the amino acid sequence of xanthine phosphoribosyl transferase polypeptide of Table 1 [SEQ ID NO:2] in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, deleted or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of xanthine phosphoribosyl transferase.

Further preferred embodiments of the invention are polynucleotides that are at least 70% identical over their entire length to a polynucleotide encoding xanthine phosphoribosyl transferase polypeptide having an amino acid sequence set out in Table 1 [SEQ ID NOS:2 and 4], and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding xanthine phosphoribosyl transferase polypeptide of the deposited strain and polynucleotides complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by the DNA of Table 1 [SEQ ID NO:1].

The invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml

denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein.

5 The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 or SEQ ID NO:3 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or a fragment thereof; and isolating said DNA sequence. Fragments
10 useful for obtaining such a polynucleotide include, for example, probes and primers described elsewhere herein.

 As discussed additionally herein regarding polynucleotide assays of the invention, for instance, polynucleotides of the invention as discussed above, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones
15 encoding xanthine phosphoribosyl transferase and to isolate cDNA and genomic clones of other genes that have a high sequence similarity to the xanthine phosphoribosyl transferase gene. Such probes generally will comprise at least 15 bases. Preferably, such probes will have at least 30 bases and may have at least 50 bases. Particularly preferred probes will have at least 30 bases and will have 50 bases or less.

20 For example, the coding region of the xanthine phosphoribosyl transferase gene may be isolated by screening using the DNA sequence provided in SEQ ID NO: 1 to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

25 The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for disease, particularly human disease, as further discussed herein relating to polynucleotide assays.

 Polynucleotides of the invention that are oligonucleotides derived from the sequences of SEQ ID NOS:1 and/or 2 may be used in the processes herein as described, but preferably for
30 PCR, to determine whether or not the polynucleotides identified herein in whole or in part are

transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

The invention also provides polynucleotides that may encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

Vectors, host cells, expression

The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold

Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

5 Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, enterococci *E. coli*, streptomyces and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293 and Bowes melanoma cells; and plant cells.

10 A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal, episomal and virus-derived vectors, *e.g.*, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses,
15 pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this
20 regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

 For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be
25 incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

 Polypeptides of the invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography,
30 hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and lectin chromatography. Most preferably, high performance liquid

chromatography is employed for purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

Diagnostic Assays

5 This invention is also related to the use of the xanthine phosphoribosyl transferase polynucleotides of the invention for use as diagnostic reagents. Detection of xanthine phosphoribosyl transferase in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of a disease. Eukaryotes (herein also "individual(s)"), particularly mammals, and especially humans, infected with an organism comprising the xanthine
10 phosphoribosyl transferase gene may be detected at the nucleic acid level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from an infected individual's cells and tissues, such as bone, blood, muscle, cartilage, and skin. Genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification technique prior
15 to analysis. RNA or cDNA may also be used in the same ways. Using amplification, characterization of the species and strain of prokaryote present in an individual, may be made by an analysis of the genotype of the prokaryote gene. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the genotype of a reference sequence. Point mutations can be identified by hybridizing amplified DNA to labeled xanthine
20 phosphoribosyl transferase polynucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in the electrophoretic mobility of the DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, e.g., Myers et al., *Science*, 230: 1242 (1985). Sequence changes at
25 specific locations also may be revealed by nuclease protection assays, such as RNase and S1 protection or a chemical cleavage method. See, e.g., Cotton et al., *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

Cells carrying mutations or polymorphisms in the gene of the invention may also be detected at the DNA level by a variety of techniques, to allow for serotyping, for example. For
30 example, RT-PCR can be used to detect mutations. It is particularly preferred to used RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA or cDNA

may also be used for the same purpose, PCR or RT-PCR. As an example, PCR primers complementary to a nucleic acid encoding xanthine phosphoribosyl transferase can be used to identify and analyze mutations. Examples of representative primers are shown below in Table 2.

5 **Table 2**

Primers for amplification of xanthine phosphoribosyl transferase polynucleotides

<u>SEQ ID NO</u>	<u>PRIMER SEQUENCE</u>
10 5	5'-TCCTC AAGGTAGATT CCTTTTAAAC-3'
6	5'-CTCCTTAAATACGACCTGAC-3'

15 The invention further provides these primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying xanthine phosphoribosyl transferase DNA isolated from a sample derived from an individual. The primers may be used to amplify the gene isolated from an infected individual such that the gene may then be subject to various techniques for elucidation of the DNA sequence. In this way, mutations in the DNA sequence may be detected and used to diagnose infection and to serotype and/or classify the infectious agent.

20 The invention further provides a process for diagnosing, disease, preferably bacterial infections, more preferably infections by *Streptococcus pneumoniae*, and most preferably otitis media, conjunctivitis, pneumonia, bacteremia, meningitis, sinusitis, pleural empyema and endocarditis, and most particularly meningitis, such as for example infection of cerebrospinal fluid, comprising determining from a sample derived from an individual a increased level of
25 expression of polynucleotide having the sequence of Table 1 [SEQ ID NO: 1]. Increased or decreased expression of xanthine phosphoribosyl transferase polynucleotide can be measured using any on of the methods well known in the art for the quantation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods.

30 In addition, a diagnostic assay in accordance with the invention for detecting over-expression of xanthine phosphoribosyl transferase protein compared to normal control tissue

samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a xanthine phosphoribosyl transferase protein, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

5 **Antibodies**

The polypeptides of the invention or variants thereof, or cells expressing them can be used as an immunogen to produce antibodies immunospecific for such polypeptides. "Antibodies" as used herein includes monoclonal and polyclonal antibodies, chimeric, single chain, simianized antibodies and humanized antibodies, as well as Fab fragments, including the
10 products of an Fab immunoglobulin expression library.

Antibodies generated against the polypeptides of the invention can be obtained by administering the polypeptides or epitope-bearing fragments, analogues or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can
15 be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic
20 mice, or other organisms such as other mammals, may be used to express humanized antibodies.

Alternatively phage display technology may be utilized to select antibody genes with binding activities towards the polypeptide either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-xanthine phosphoribosyl transferase or from naive libraries (McCafferty, J. *et al.*, (1990), *Nature* 348, 552-554; Marks, J. *et al.*, (1992)
25 *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by chain shuffling (Clackson, T. *et al.*, (1991) *Nature* 352, 624-628).

If two antigen binding domains are present each domain may be directed against a different epitope - termed 'bispecific' antibodies.

The above-described antibodies may be employed to isolate or to identify clones
30 expressing the polypeptides to purify the polypeptides by affinity chromatography.

Thus, among others, antibodies against xanthine phosphoribosyl transferase- polypeptide may be employed to treat infections, particularly bacterial infections and especially otitis media, conjunctivitis, pneumonia, bacteremia, meningitis, sinusitis, pleural empyema and endocarditis, and most particularly meningitis, such as for example infection of cerebrospinal fluid.

5 Polypeptide variants include antigenically, epitopically or immunologically equivalent variants that form a particular aspect of this invention. The term "antigenically equivalent derivative" as used herein encompasses a polypeptide or its equivalent which will be specifically recognized by certain antibodies which, when raised to the protein or polypeptide according to the invention, interfere with the immediate physical interaction between pathogen
10 and mammalian host. The term "immunologically equivalent derivative" as used herein encompasses a peptide or its equivalent which when used in a suitable formulation to raise antibodies in a vertebrate, the antibodies act to interfere with the immediate physical interaction between pathogen and mammalian host.

The polypeptide, such as an antigenically or immunologically equivalent derivative or
15 a fusion protein thereof is used as an antigen to immunize a mouse or other animal such as a rat or chicken. The fusion protein may provide stability to the polypeptide. The antigen may be associated, for example by conjugation, with an immunogenic carrier protein for example bovine serum albumin (BSA) or keyhole limpet haemocyanin (KLH). Alternatively a multiple antigenic peptide comprising multiple copies of the protein or polypeptide, or an antigenically
20 or immunologically equivalent polypeptide thereof may be sufficiently antigenic to improve immunogenicity so as to obviate the use of a carrier.

Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized"; where the complementarity determining region(s) of the hybridoma-derived
25 antibody has been transplanted into a human monoclonal antibody, for example as described in Jones, P. et al. (1986), *Nature* 321, 522-525 or Tempest et al., (1991) *Biotechnology* 9, 266-273.

The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles
30 (Wolff et al., *Hum Mol Genet* 1992, 1:363, Manthorpe et al., *Hum. Gene Ther.* 1993:4, 419), delivery of DNA complexed with specific protein carriers (Wu et al., *J Biol Chem.* 1989:

264,16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, 1986:83,9551), encapsulation of DNA in various forms of liposomes (Kaneda et al., *Science* 1989:243,375), particle bombardment (Tang et al., *Nature* 1992, 356:152, Eisenbraun et al., *DNA Cell Biol* 1993, 12:791) and *in vivo* infection using cloned retroviral vectors (Seeger et al., *PNAS USA* 1984:81,5849).

Antagonists and agonists - assays and molecules

Polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, *e.g.*, Coligan et al., *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of xanthine phosphoribosyl transferase polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bacteriocidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising xanthine phosphoribosyl transferase polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a xanthine phosphoribosyl transferase agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the xanthine phosphoribosyl transferase polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of xanthine phosphoribosyl transferase polypeptide are most likely to be good antagonists. Molecules that bind well and increase the rate of product production from substrate are agonists. Detection of the rate or level of production of product from substrate may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric labeled substrate converted into product, a reporter gene that is responsive to changes in xanthine phosphoribosyl transferase polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for xanthine phosphoribosyl transferase antagonists is a competitive assay that combines xanthine phosphoribosyl transferase and a potential antagonist

with xanthine phosphoribosyl transferase-binding molecules, recombinant xanthine phosphoribosyl transferase binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. Xanthine phosphoribosyl transferase can be labeled, such as by radioactivity or a colorimetric compound, such that the number of xanthine phosphoribosyl transferase molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide or polypeptide of the invention and thereby inhibit or extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing xanthine phosphoribosyl transferase-induced activities, thereby preventing the action of xanthine phosphoribosyl transferase by excluding xanthine phosphoribosyl transferase from binding.

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of xanthine phosphoribosyl transferase.

Each of the DNA sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the DNA sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide or inhibitor of the invention to interfere with the initial physical interaction between a pathogen and mammalian host responsible for sequelae of infection. In particular the molecules of the

invention may be used: in the prevention of adhesion of bacteria, in particular gram positive bacteria, to mammalian extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block xanthine phosphoribosyl transferase protein-mediated mammalian cell invasion by, for example, initiating phosphorylation of mammalian tyrosine
5 kinases (Rosenshine *et al.*, *Infect. Immun.* 60:2211 (1992); to block bacterial adhesion between mammalian extracellular matrix proteins and bacterial xanthine phosphoribosyl transferase proteins that mediate tissue damage and; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

10 The antagonists and agonists of the invention may be employed, for instance, to inhibit and treat otitis media, conjunctivitis, pneumonia, bacteremia, meningitis, sinusitis, pleural empyema and endocarditis, and most particularly meningitis, such as for example infection of cerebrospinal fluid.

Vaccines

15 Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal which comprises inoculating the individual with xanthine phosphoribosyl transferase, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Streptococcus pneumoniae* infection. Also provided
20 are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector to direct expression of xanthine phosphoribosyl transferase, or a fragment or a variant thereof, for expressing xanthine phosphoribosyl transferase, or a fragment or a variant thereof *in vivo* in
25 order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual from disease, whether that disease is already established within the individual or not. One way of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise.

30 Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid.

A further aspect of the invention relates to an immunological composition which, when introduced into an individual capable or having induced within it an immunological response, induces an immunological response in such individual to a xanthine phosphoribosyl transferase or protein coded therefrom, wherein the composition comprises a recombinant xanthine phosphoribosyl transferase or protein coded therefrom comprising DNA which codes for and expresses an antigen of said xanthine phosphoribosyl transferase or protein coded therefrom. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity or cellular immunity such as that arising from CTL or CD4+ T cells.

A xanthine phosphoribosyl transferase polypeptide or a fragment thereof may be fused with co-protein which may not by itself produce antibodies, but is capable of stabilizing the first protein and producing a fused protein which will have immunogenic and protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Hemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, relatively large co-proteins which solubilize the protein and facilitate production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system. The co-protein may be attached to either the amino or carboxy terminus of the first protein.

Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof which have been shown to encode non-variable regions of bacterial cell surface proteins in DNA constructs used in such genetic immunization experiments in animal models of infection with *Streptococcus pneumoniae* will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value from the requisite organ of the animal successfully resisting or clearing infection for the development of prophylactic agents or therapeutic

treatments of bacterial infection, particularly *Streptococcus pneumoniae* infection, in mammals, particularly humans.

5 The polypeptide may be used as an antigen for vaccination of a host to produce specific antibodies which protect against invasion of bacteria, for example by blocking adherence of bacteria to damaged tissue. Examples of tissue damage include wounds in skin or connective tissue caused, e.g., by mechanical, chemical or thermal damage or by implantation of indwelling devices, or wounds in the mucous membranes, such as the mouth, mammary glands, urethra or vagina.

10 The invention also includes a vaccine formulation which comprises an immunogenic recombinant protein of the invention together with a suitable carrier. Since the protein may be broken down in the stomach, it is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the
15 formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may
20 also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

25 While the invention has been described with reference to certain xanthine phosphoribosyl transferase protein, it is to be understood that this covers fragments of the naturally occurring protein and similar proteins with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant protein.

Compositions, kits and administration

30 The invention also relates to compositions comprising the polynucleotide or the polypeptides discussed above or their agonists or antagonists. The polypeptides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to a subject.

Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

Alternatively the composition may be formulated for topical application for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

In-dwelling devices include surgical implants, prosthetic devices and catheters, i.e., devices that are introduced to the body of an individual and remain in position for an extended time. Such devices include, for example, artificial joints, heart valves, pacemakers, vascular grafts, vascular catheters, cerebrospinal fluid shunts, urinary catheters, continuous ambulatory
5 peritoneal dialysis (CAPD) catheters.

The composition of the invention may be administered by injection to achieve a systemic effect against relevant bacteria shortly before insertion of an in-dwelling device. Treatment may be continued after surgery during the in-body time of the device. In addition, the composition could also be used to broaden perioperative cover for any surgical technique to
10 prevent bacterial wound infections, especially *Streptococcus pneumoniae* wound infections.

Many orthopaedic surgeons consider that humans with prosthetic joints should be considered for antibiotic prophylaxis before dental treatment that could produce a bacteremia. Late deep infection is a serious complication sometimes leading to loss of the prosthetic joint and is accompanied by significant morbidity and mortality. It may therefore be possible to
15 extend the use of the active agent as a replacement for prophylactic antibiotics in this situation.

In addition to the therapy described above, the compositions of this invention may be used generally as a wound treatment agent to prevent adhesion of bacteria to matrix proteins exposed in wound tissue and for prophylactic use in dental treatment as an alternative to, or in conjunction with, antibiotic prophylaxis.

20 Alternatively, the composition of the invention may be used to bathe an indwelling device immediately before insertion. The active agent will preferably be present at a concentration of 1µg/ml to 10mg/ml for bathing of wounds or indwelling devices.

A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5
25 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

Each reference disclosed herein is incorporated by reference herein in its entirety. Any
30 patent application to which this application claims priority is also incorporated by reference herein in its entirety.

EXAMPLES

5 The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1 Strain selection, Library Production and Sequencing

10 The polynucleotide having the DNA sequence given in SEQ ID NO:1 was obtained from a library of clones of chromosomal DNA of *Streptococcus pneumoniae* in *E. coli*. The sequencing data from two or more clones containing overlapping *Streptococcus pneumoniae* DNAs was used to construct the contiguous DNA sequence in SEQ ID NO:1. Libraries may be prepared by routine methods, for example:

Methods 1 and 2 below.

15 Total cellular DNA is isolated from *Streptococcus pneumoniae* 0100993 according to standard procedures and size-fractionated by either of two methods.

Method 1

20 Total cellular DNA is mechanically sheared by passage through a needle in order to size-fractionate according to standard procedures. DNA fragments of up to 11kbp in size are rendered blunt by treatment with exonuclease and DNA polymerase, and EcoRI linkers added. Fragments are ligated into the vector Lambda ZapII that has been cut with EcoRI, the library packaged by standard procedures and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

Method 2

25 Total cellular DNA is partially hydrolyzed with a one or a combination of restriction enzymes appropriate to generate a series of fragments for cloning into library vectors (e.g., RsaI, PstI, AluI, BshI235I), and such fragments are size-fractionated according to standard procedures. EcoRI linkers are ligated to the DNA and the fragments then ligated into the vector Lambda ZapII that have been cut with EcoRI, the library packaged by standard procedures, and *E.coli* infected with the packaged library. The library is amplified by standard
30 procedures.

SEQUENCE LISTING

5 **(1) GENERAL INFORMATION**

(i) APPLICANT: Burnham, Martin K.

10 **(ii) TITLE OF THE INVENTION: NOVEL XANTHINE PHOSPHORIBOSYL
 TRANSFERASE**

(iii) NUMBER OF SEQUENCES: 6

(iv) CORRESPONDENCE ADDRESS:

15 **(A) ADDRESSEE: Dechert Price & Rhoads**

(B) STREET: 997 Lenox Drive, Building 3, Suite 210

(C) CITY: Lawrenceville

(D) STATE: NJ

(E) COUNTRY: USA

20 **(F) ZIP: 08543**

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette

(B) COMPUTER: IBM Compatible

25 **(C) OPERATING SYSTEM: DOS**

(D) SOFTWARE: FastSEQ for Windows Version 2.0

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

30 **(B) FILING DATE:**

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

5

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Bloom, Allen

10 **(B) REGISTRATION NUMBER: 29,135**

(C) REFERENCE/DOCKET NUMBER:

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 609-520-3214

15 **(B) TELEFAX: 609-520-3259**

(C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 582 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

25 **(D) TOPOLOGY: linear**

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

30 ATGAAATTAT TAGAAGAGCG CATCCTCAAG GATGGGCATA TCTTGGGTGA
TAACATCCTC 60

AAGGTAGATT CCTTTTAAAC CCACCAAGTT GACTTTAGCT TGATGCGAGA
GATTGGTAAG 120

GTTTTTGCGG AAAAATTTGC TGCTACTGGC ATTACCAAGG TCGTAACCAT
TGAAGCGTCG 180

5 GGTATTGCCC CAGCCGTTTT TACAGCTGAA GCCTTAAACG TTCCCATGAT
TTTCGCCAAA 240

AAAGCTAAGA ACATCACCAT GAACGAAGGC ATCTTAACTG CTCAAGTCTA
CTCCTTTACC 300

AAGCAGGTGA CCAGCACTGT TTCTATCGCT GGAAAATTCC TCTCACCAGA
10 GGACAAGGTT 360

TTGATTATCG ACGATTTTCCT TGCTAATGGC CAAGCTGCTA AAGGCTTGAT
TCAAATCATC 420

GAACAGGCCG GTGCCACAGT CCAAGCTATC GGTATCGTGA TTGAGAAATC
CTTCCAAGAT 480

15 GGTCGTGATT TGCTTGAAAA AGCAGGCTAC CCTGTCCTAT CACTTGCTCG
CTTGGATCGT 540

TTTGAAAATG GTCAGGTCGT ATTTAAGGAG GCAGATCTCT AA
582

20 (2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 193 amino acids

(B) TYPE: amino acid

25 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

30

Met Lys Leu Leu Glu Glu Arg Ile Leu Lys Asp Gly His Ile Leu Gly

1 5 10 15
 Asp Asn Ile Leu Lys Val Asp Ser Phe Leu Thr His Gln Val Asp Phe
 20 25 30
 Ser Leu Met Arg Glu Ile Gly Lys Val Phe Ala Glu Lys Phe Ala Ala
 5 35 40 45
 Thr Gly Ile Thr Lys Val Val Thr Ile Glu Ala Ser Gly Ile Ala Pro
 50 55 60
 Ala Val Phe Thr Ala Glu Ala Leu Asn Val Pro Met Ile Phe Ala Lys
 65 70 75 80
 10 Lys Ala Lys Asn Ile Thr Met Asn Glu Gly Ile Leu Thr Ala Gln Val
 85 90 95
 Tyr Ser Phe Thr Lys Gln Val Thr Ser Thr Val Ser Ile Ala Gly Lys
 100 105 110
 Phe Leu Ser Pro Glu Asp Lys Val Leu Ile Ile Asp Asp Phe Leu Ala
 15 115 120 125
 Asn Gly Gln Ala Ala Lys Gly Leu Ile Gln Ile Ile Glu Gln Ala Gly
 130 135 140
 Ala Thr Val Gln Ala Ile Gly Ile Val Ile Glu Lys Ser Phe Gln Asp
 145 150 155 160
 20 Gly Arg Asp Leu Leu Glu Lys Ala Gly Tyr Pro Val Leu Ser Leu Ala
 165 170 175
 Arg Leu Asp Arg Phe Glu Asn Gly Gln Val Val Phe Lys Glu Ala Asp
 180 185 190
 Leu

25

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

30

(A) LENGTH: 579 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATGAAATTAT TAGAAGAGCG CATCCTCAAG GATGGGCATA TCTTGGGTGA
TAACATCCTC 60
AAGGTAGATT CCTTTTAAAC CCACCAAGTT GACTTTAGCT TGATGCGAGA
10 GATTGGTAAG 120
GTTTTTGCGG AAAAATTGTC TGCTACTGGC ATTACCAAGG TCGTAACCAT
TGAAGCGTCG 180
GGTATTGCCC CAGCCGTTTT TACAGCTGAA GCCTTAAACG TTCCCATGAT
TTTCGCCAAA 240
15 AAAGCTAAGA ACATCACCAT GAACGAAGGC ATCTTAACTG CTCAAGTCTA
CTCCTTTACC 300
AAGCAGGTGA CCAGCACTGT TTCTATCGCT GGAAAATTCC TCTCACCAGA
GGACAAGGTT 360
TTGATTATCG ACGATTTTCCT TGCTAATGGC CAAGCTGCTA AAGGCTTGAT
20 TCAAATCATC 420
GAACAGGCCG GTGCCACAGT CCAAGCTATC GGTATCGTGA TTGAGAAATC
CTTCCAAGAT 480
GGTCGTGATT TGCTTGAAAA AGCAGGCTAC CCTGTCCTAT CACTTGCTCG
CTTGGATCGT 540
25 TTTGAAAATG GTCAGGTCGT ATTTAAGGAG GCAGATCTC 579

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 193 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Lys Leu Leu Glu Glu Arg Ile Leu Lys Asp Gly His Ile Leu Gly
1 5 10 15
Asp Asn Ile Leu Lys Val Asp Ser Phe Leu Thr His Gln Val Asp Phe
10 20 25 30
Ser Leu Met Arg Glu Ile Gly Lys Val Phe Ala Glu Lys Phe Ala Ala
35 40 45
Thr Gly Ile Thr Lys Val Val Thr Ile Glu Ala Ser Gly Ile Ala Pro
50 55 60
15 Ala Val Phe Thr Ala Glu Ala Leu Asn Val Pro Met Ile Phe Ala Lys
65 70 75 80
Lys Ala Lys Asn Ile Thr Met Asn Glu Gly Ile Leu Thr Ala Gln Val
85 90 95
Tyr Ser Phe Thr Lys Gln Val Thr Ser Thr Val Ser Ile Ala Gly Lys
20 100 105 110
Phe Leu Ser Pro Glu Asp Lys Val Leu Ile Ile Asp Asp Phe Leu Ala
115 120 125
Asn Gly Gln Ala Ala Lys Gly Leu Ile Gln Ile Ile Glu Gln Ala Gly
130 135 140
25 Ala Thr Val Gln Ala Ile Gly Ile Val Ile Glu Lys Ser Phe Gln Asp
145 150 155 160
Gly Arg Asp Leu Leu Glu Lys Ala Gly Tyr Pro Val Leu Ser Leu Ala
165 170 175
Arg Leu Asp Arg Phe Glu Asn Gly Gln Val Val Phe Lys Glu Ala Asp
30 180 185 190
Leu

(2) INFORMATION FOR SEQ ID NO:5:

5 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TCCTCAAGGT AGATTCCTTT TTAAC

25

15

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CTCCTTAAAT ACGACCTGAC

20

What is claimed is:

1. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:
 - (a) a polynucleotide having at least a 70% identity to a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:2;
 - (b) a polynucleotide having at least a 70% identity to a polynucleotide encoding the same mature polypeptide expressed by the xanthine phosphoribosyl transferase gene contained in the *Streptococcus pneumoniae* of the deposited strain;
 - (c) a polynucleotide encoding a polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2;
 - (d) a polynucleotide which is complementary to the polynucleotide of (a), (b) or (c); and
 - (e) a polynucleotide comprising at least 15 sequential bases of the polynucleotide of (a), (b), (c) or (d).
2. The polynucleotide of Claim 1 wherein the polynucleotide is DNA.
3. The polynucleotide of Claim 1 wherein the polynucleotide is RNA.
4. The polynucleotide of Claim 2 comprising the nucleic acid sequence set forth in SEQ ID NO:1.
5. The polynucleotide of Claim 2 comprising nucleotide 1 to 579 set forth in SEQ ID NO:1.
6. The polynucleotide of Claim 2 which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2.
7. A vector comprising the polynucleotide of Claim 1.
8. A host cell comprising the vector of Claim 7.
9. A process for producing a polypeptide comprising: expressing from the host cell of Claim 8 a polypeptide encoded by said DNA.
10. A process for producing a xanthine phosphoribosyl transferase polypeptide or fragment comprising culturing a host of claim 8 under conditions sufficient for the production of said polypeptide or fragment.
11. A polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2.

12. A polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2.
13. An antibody against the polypeptide of claim 11.
14. An antagonist which inhibits the activity or expression of the polypeptide of claim 11.
15. A method for the treatment of an individual in need of xanthine phosphoribosyl transferase polypeptide comprising: administering to the individual a therapeutically effective amount of the polypeptide of claim 11.
16. A method for the treatment of an individual having need to inhibit xanthine phosphoribosyl transferase polypeptide comprising: administering to the individual a therapeutically effective amount of the antagonist of Claim 14.
17. A process for diagnosing a disease related to expression or activity of the polypeptide of claim 11 in an individual comprising:
 - (a) determining a nucleic acid sequence encoding said polypeptide, and/or
 - (b) analyzing for the presence or amount of said polypeptide in a sample derived from the individual.
18. A method for identifying compounds which interact with and inhibit or activate an activity of the polypeptide of claim 11 comprising:

contacting a composition comprising the polypeptide with the compound to be screened under conditions to permit interaction between the compound and the polypeptide to assess the interaction of a compound, such interaction being associated with a second component capable of providing a detectable signal in response to the interaction of the polypeptide with the compound;

and determining whether the compound interacts with and activates or inhibits an activity of the polypeptide by detecting the presence or absence of a signal generated from the interaction of the compound with the polypeptide.
19. A method for inducing an immunological response in a mammal which comprises inoculating the mammal with xanthine phosphoribosyl transferase polypeptide of claim 11, or a fragment or variant thereof, adequate to produce antibody and/or T cell immune response to protect said animal from disease.
20. A method of inducing immunological response in a mammal which comprises delivering a nucleic acid vector to direct expression of xanthine phosphoribosyl transferase polypeptide of claim 11, or fragment or a variant thereof, for expressing said xanthine

phosphoribosyl transferase polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response to produce antibody and/ or T cell immune response to protect said animal from disease.

ABSTRACT OF THE DISCLOSURE

The invention provides xanthine phosphoribosyl transferase polypeptides and DNA (RNA) encoding xanthine phosphoribosyl transferase polypeptides and methods for producing such polypeptides by recombinant techniques. Also provided are methods for utilizing xanthine phosphoribosyl transferase polypeptides to screen for antibacterial compounds.